

CHAPTER 9

ALASKA PLAICE

by

Thomas K. Wilderbuer, Daniel G. Nichol, and Paul D. Spencer

Executive Summary

The following changes have been made to this assessment relative to the November 2005 SAFE:

Changes in the assessment input data

- 1) The 2006 catch data was updated, and catch through 8 September, 2007 were included in the assessment.
- 2) The 2007 trawl survey biomass estimate and standard error, and the 2007 survey length composition were included in the assessment.
- 3) The 2006 survey ages were read and the 2006 survey age composition was added to the assessment.
- 4) Length bins and the transition matrix were extended from 45 to 60 cm.

Model results

- 1) Estimated 3+ total biomass for 2008 is 1,854,000 t.
- 2) Projected female spawning biomass for 2008 is 335,900 t.
- 3) Recommended ABC for 2008 is 194,100 t based on an $F_{40\%} = 0.59$ harvest level.
- 4) 2008 overfishing level is 247,500 t based on a $F_{35\%}$ (0.81) harvest level.

	2007 Assessment recommendations for the 2008 harvest	2006 Assessment recommendations for the 2007 harvest
ABC	194,100 t	189,900 t
Overfishing	247,500 t	241,200 t
F_{ABC}	$F_{0.40} = 0.59$	$F_{0.40} = 0.61$
$F_{\text{overfishing}}$	$F_{0.35} = 0.81$	$F_{0.35} = 0.83$
Projected total biomass	1,854,000 t	1,335,200 t
Projected fem. spawning biomass	335,900 t	294,800 t

SSC Comments from December 2005

Many of the length frequency plots show very large proportions of fish in the largest length class. The SSC requests that the authors consider extending the range of length bins to better mimic the dynamics of the larger fish.

The length bins and the transition matrix were extended from 45 to 60 cm to improve the population modeling of larger fish.

While the assessment takes account of differences in weight at age between sexes when computing biomass, the SSC recommends that the assessment author consider moving to a fully split-sex model. Such a model would allow differing dynamics beyond the age of maturation to be captured more fully

This is an excellent suggestion which the lead author will try to implement before the next assessment cycle.

Introduction

Prior to 2002, Alaska plaice (*Pleuronectes quadrituberculatus*) were managed as part of the “other flatfish” complex. Since then an age-structured model has been used for the stock assessment allowing Alaska plaice to be managed separately from the “other flatfish” complex as a single species.

The distribution of Alaska plaice is mainly on the Eastern Bering Sea continental shelf, with only small amounts found in the Aleutian Islands region. In particular, the summer distribution of Alaska plaice is generally confined to depths < 110 m, with larger fish predominately in deep waters and smaller juveniles (<20 cm) in shallow coastal waters (Zhang et al., 1998). The Alaska plaice distribution overlaps with rock sole (*Lepidopsetta polyxystra*) and yellowfin sole (*Limanda aspera*), but the center of the distribution is north of the center of the other two species.

Catch History

Catches of Alaska plaice increased from approximately 1,000 t in 1971 to a peak of 62,000 t in 1988, the first year of joint venture processing (JVP) (Table 9.1). Part of this apparent increase was due to increased species identification and reporting of catches in the 1970s. Because of the overlap of the Alaska plaice distribution with that of yellowfin sole, much of the Alaska plaice catch during the 1960s was likely caught as bycatch in the yellowfin sole fishery (Zhang et al. 1998). With the cessation of joint venture fishing operations in 1991, Alaska plaice are now harvested exclusively by domestic vessels. Catch data from 1980-89 by its component fisheries (JVP, non-U.S., and domestic) are available in Wilderbuer and Walters (1990). The catch of Alaska plaice taken in research surveys from 1977 –2007 are shown in Table 9.2.

Since implementation of the Magnuson Fishery Conservation and Management Act (MFCMA) in 1977, Alaska plaice generally have been lightly harvested as no major commercial target fishery exists for them. The 2007 catch (through 8 September) was 19,176 t, primarily caught in pursuit of other flatfish species. Alaska plaice are grouped with the rock sole, flathead sole, and other flatfish fisheries under a common prohibited species catch (PSC) limit, with seasonal and total annual allowances of prohibited species bycatch by these flatfish fisheries applied to the fisheries within the group. In recent years, these fisheries have been closed prior to attainment of the TAC due to the bycatch of halibut (Table 9.3), and typically are also closed during the first quarter due to a seasonal bycatch cap. Alaska plaice were placed on bycatch status each spring of the past three years due to the attainment of a very low TAC (relative to the ABC) for this species.

Substantial amounts of Alaska plaice are discarded in various eastern Bering Sea target fisheries due to the low market interest. Retained and discarded catches were reported for Alaska plaice for the first time in 2002, and indicated that of the 12,176 t caught only 370 t were retained, resulting in a retention rate of 3.0 % (Table 9.4). Similar patterns were observed for 2003 - 2005 (4%, 5% and 6%, respectively). The amount of Alaska plaice retained in 2006 improved to 15%. Examination of the discard data, by fishery, indicates that 81% - 87% of the discards in 2002 - 2006 can be attributed to the yellowfin sole fishery. Discarding also occurred in the rock sole,

flathead sole, and Pacific cod fisheries. The locations where Alaska plaice were caught, by month, in 2007 are shown in Figure 9.1.

Data

Fishery Catch and Catch-at-Age Data

This assessment uses fishery catches from 1971 through 8 September, 2007 (Table 9.2). Fishery length compositions from 1975-76, 1978-89, 1993, 1995, and 2001 were also used, as well as age compositions from 2000, 2002 and 2003. The number of ages and lengths sampled from the fishery are shown in Table 9.5.

Survey Data

Because Alaska plaice are usually taken incidentally in target fisheries for other species, CPUE from commercial fisheries is considered unreliable information for determining trends in abundance for these species. It is therefore necessary to use research vessel survey data to assess the condition of these stocks.

Large-scale bottom trawl surveys of the Eastern Bering Sea continental shelf have been conducted in 1975 and 1979-2007 by NMFS. Survey estimates of total biomass and numbers at age are shown in Tables 9.6 and 9.7, respectively. It should be recognized that the resultant biomass estimates are point estimates from an "area-swept" survey. As a result, they carry the uncertainty inherent in the technique. It is assumed that the sampling plan covers the distribution of the fish and that all fish in the path of the trawl are captured. That is, there are no losses due to escape or gains due to gear herding effects. Trawl survey estimates of Alaska plaice biomass increased dramatically from 1975 through 1982 and have remained at a high and stable level since (Table 9.6, Figure 9.2).

The trawl gear was changed in 1982 from the 400 mesh eastern trawl to the 83-112 trawl, as the latter trawl has better bottom contact. This may contribute to the increase in Alaska plaice seen from 1981 to 1982, as increases between these years were noticed in other flatfish as well. However, large changes in Alaska plaice biomass between adjacent years have occurred without changes in trawl gear, such as the increase from 1980 to 1981 and the decrease from 1984 to 1985.

Although calibration between years with different trawl gear has not been accomplished, the survey data since 1982 does incorporate calibration between the two vessels used in the survey. Fishing Power Coefficients (FPC) were estimated following the methods of Kappenman (1992). The trend of the biomass estimates is the same as without the calibration between vessels, but the magnitude of the change in 1988 was markedly reduced. In 1988, one vessel had slightly smaller and lighter trawl doors which may have affected the estimates for several species. With the exception of the 1988 estimate, Alaska plaice has shown a relatively stable trend since 1985, although abundance was higher in the 1994, 1997 and 2006 surveys. The 2007 estimate of 421,765 t is similar to the survey estimates between 1998 and 2005.

Assessments for other BSAI flatfish have suggested a relationship between bottom temperature and survey catchability (Wilderbuer et al. 2002), where bottom temperatures are hypothesized to affect survey catchability by affecting either stock distributions and/or the activity level of flatfish relative to the capture process. This relationship was investigated for Alaska plaice by using the annual temperature anomalies from surveys conducted from 1982 to 2004. Much of the trend in survey biomass estimates of Alaska plaice is expected to be explained by changes in stock biomass rather than survey catchability, and this trend was fit with a LOWESS smoother. The residuals from the smoothed trend produce a detrended estimate of survey biomass, which was then standardized and compared to the bottom temperature anomalies (Figure 9.3). Little correspondence exists between the two time series, and the cross-correlation coefficient (-0.17) was not significant at the 0.05 level. Thus, the relationship between bottom temperature and survey catchability was not pursued further.

Survey Length, Weight and Age Information

In previous assessments, information regarding growth of Alaska plaice was produced by fitting a von Bertalanffy curve to the available length-at-age data from specimens sampled in trawl surveys. However, such data are typically obtained from length-stratified sampling, thus potentially introducing some bias into estimates of length at age (Kimura and Chikuni 1987). In this assessment, the estimated population numbers at length was multiplied by the age-length key in order to produce a matrix of estimated population numbers by age and length, from which an unbiased average length for each age can be determined. Because separate length-stratified samples of otoliths occur for the northwest and southeast EBS shelf, this procedure was conducted separately in each area, and a single average length at age was obtained by taking an average of the two estimates (weighted by population size). Separate growth curves were produced for each year where aged otoliths were available, which includes 1982, 1988, 1992-1995, 1998, and 2000-2002. The number of age and length samples obtained from the surveys are shown in Table 9.8.

With the exception of age 5, consistent temporal trends in the mean length at age were not observed (Figure 9.4), suggesting that a single growth curve over all modeled years can suitably represent the pattern in length at age. The von Bertalanffy parameters were estimated as:

$L_{inf}(cm)$	k	t_0
45.6	0.1315	0.1334

Note that these estimates are similar to those estimated in the 2003 assessment, which were $L_{inf} = 47.0$, $k = 0.1269$, and $t_0 = -0.57$. The length-weight relationship of the form $W = aL^b$ was also updated from the available data, with parameter estimates of $a = 0.007$ and $b = 3.15$ obtained from the 2001-2002 survey data. The combination of the weight-length relationship and the von Bertalanffy growth curve produces an estimated weight-at-age relationship that is similar to that used in previous Alaska plaice assessments (Figure 9.5).

In summary, the data available for Alaska plaice are

-
- 1) Total catch weight, 1971-2007;
 - 2) Proportional catch number at age, 2000,2002-2003
 - 3) Proportional catch number at length, 1975-76, 1978-89, 1993, 1995, 2000
 - 4) Survey biomass and standard error 1975, 1979-2007;
 - 5) Survey age composition, 1982, 1988, 1992-1995, 1998, 2000-2002, 2005, 2006
 - 6) Survey length composition, 1983-1987, 1989-1991, 1996-1997, 1999, 2003, 2004, 2007
-

Analytical Approach

Model Structure

A catch-at-age population dynamics model was used to obtain estimates of several population variables of the Alaska plaice stock, including recruitment, population size, and catch. This catch at age model was developed with the software program AD Modelbuilder. Population size in numbers at age a in year t was modeled as

$$N_{t,a} = N_{t-1,a-1} e^{-Z_{t-1,a-1}} \quad 3 \leq a < A, \quad 3 \leq t \leq T$$

where Z is the sum of the instantaneous fishing mortality rate ($F_{t,a}$) and the natural mortality rate (M), A is the maximum modeled age in the population, and T is the terminal year of the analysis. The numbers at age A are a “pooled” group consisting of fish of age A and older, and are estimated as

$$N_{t,A} = N_{t-1,A-1} e^{-Z_{t-1,A-1}} + N_{t-1,A} e^{-Z_{t-1,A}}$$

Recruitment was modeled as the number of age 3 fish. The efficacy of estimating productivity directly from the stock-recruitment data (as opposed to using an SPR proxy) was examined by comparing results from fitting either the Ricker or Beverton-Holt forms within the model, and is described in more detail in the “Tier 1 evaluation” section below. Briefly, recruits were modeled as

$$R_t = f(S_{t-a_r}) e^{v_t}$$

where R is age 3 recruits, $f(S)$ is the form of the stock-recruitment function, S is spawning stock size, v is random error, and a_r is the age of recruitment.

The numbers at age in the first year are modeled with a lognormal distribution

$$N_{1,a} = e^{(meaninit - M(a-1) + \gamma_a)}$$

where *meaninit* is the mean and γ is an age-variant deviation.

The mean numbers at age within each year were computed as

$$\bar{N}_{t,a} = N_{t,a} * (1 - e^{-Z_{t,a}}) / Z_{t,a}$$

Catch in numbers at age in year t ($C_{t,a}$) and total biomass of catch each year were modeled as

$$C_{t,a} = F_{t,a} \bar{N}_{t,a}$$

$$Y_t = \sum_{a=1}^A C_{t,a} w_a$$

where w_a is the mean weight at age for plaice.

A transition matrix was derived from the von Bertalanffy growth relationship, and used to convert the modeled numbers at age into modeled numbers at length. There are 36 length bins ranging from 10 to 45 cm, and 23 age groups ranging from 3 to 25+. For each modeled age, the transition matrix consists of a probability distribution of numbers at length, with the expected value equal to the predicted length-at-age from the von Bertalanffy relationship. The variation around this expected value was derived from a linear regression of coefficient of variation (CV) in length-at-age against age, where the CV were obtained from the sampled specimens over all survey years. The estimated linear relationship predicts a CV of 0.14 at age 3 and a CV of 0.10 at age 25. The transition matrix, vector of mean numbers at age, and survey selectivity by age were used to compute the estimated survey length composition, by year, as

$$\bar{\mathbf{N}}_t = (\mathbf{srvsel} * \bar{\mathbf{N}}_t) * \mathbf{TR}^T$$

where \mathbf{srvsel} is a vector of survey selectivity by age.

Estimating certain parameters in different stages enhances the estimation of large number of parameters in nonlinear models. For example, the fishing mortality rate for a specific age and time ($F_{t,a}$) is modeled as the product of an age-specific selectivity function ($fishsel_a$) and a year-specific fully-selected fishing mortality rate. The fully selected mortality rate is modeled as the product of a mean (μ) and a year-specific deviation (ε_t), thus $F_{t,a}$ is

$$F_{t,a} = fishsel_a * e^{(\mu + \varepsilon_t)}$$

In the early stages of parameter estimation, the selectivity coefficients are not estimated. As the solution is being approached, selectivity was modeled with the logistic function:

$$fishsel_a = \frac{1}{1 + e^{(-slope(a - fifty))}}$$

where the parameter $slope$ affects the steepness of the curve and the parameter $fifty$ is the age at which sel_a equals 0.5. The selectivity for the survey is modeled in a similar manner.

Estimation of maximum sustainable yield

F_{msy} for Alaska plaice was estimated using the Ricker and Beverton-Holt stock recruitment curves. Additionally, for each type of curve we make separate estimates of F_{msy} based upon all year classes available or the post-1989 year classes, corresponding to differing hypotheses regarding “regime shifts”. The two different forms of recruitment curves were used because they correspond to differing assumptions regarding the nature of density-dependence in the early life-history period. For example, the strongly density dependent patterns possible in the Ricker curve may be caused by cannibalism, the transmission of disease, or density-dependent growth coupled with size-dependant predation. Alternatively, mechanisms such as competition for food or space correspond to the Beverton-Holt model (Hilborn and Walters 1992).

Briefly, a stock recruitment curve is fit to the available data, from which an equilibrium level of recruitment is solved for each level of fishing mortality. A yield curve (identifying equilibrium yield as a function of fishing mortality) is generated by multiplying equilibrium recruitment by yield per recruit, where each term in this product is a function of fishing mortality. The maximum sustainable yield is identified as the point where the derivative of the yield curve is zero, and the fishing mortality associated with MSY is F_{msy} .

The function form used for the Ricker stock recruitment curve was

$$R = \alpha S e^{-\beta S}$$

and the Beverton-Holt functional form was

$$R = \frac{\alpha S}{\beta + S}$$

where α and β are parameters corresponding to density-dependent and density-independent processes, respectively. A convenient reparameterization expresses the original stock-recruitment curve as function of R_0 (the recruitment associated with an unfished stock, or S_0) and the dimensionless steepness parameter h (the proportion of R_0 attained when the stock size is 20% of S_0). Note that for the Beverton-Holt curve, this scales the slope at the origin of the stock-recruitment curve into the interval (0.2,1.0). For the Ricker curve, this reparameterization is achieved by the following substitutions for α and β :

$$\alpha = \frac{(5h)^{\frac{5}{4}}}{\phi} \quad \text{and} \quad \beta = \frac{5 \ln(5h)}{4\phi R_0}$$

where ϕ is the spawner-per-recruit associated with no fishing, which is a constant dependent upon the size at age, proportion mature at age, and natural mortality. For the Beverton-Holt curve, the following substitution is required for the reparameterization:

$$\alpha = \frac{0.8R_0 h}{h - 0.2} \quad \text{and} \quad \beta = \frac{0.2\phi R_0 (1 - h)}{(h - 0.2)}$$

The equilibrium recruitment, at a particular level of fishing mortality, for the Ricker curve is

$$R_{eq} = \frac{-\ln\left(\frac{1}{\alpha\phi}\right)}{\phi\beta}$$

where ϕ is the spawner per recruit associated with a particular level of fishing mortality, and is a function of size at age, proportion mature at age, fishing selectivity, and fishing and natural mortality.

For the Beverton-Holt curve, the equilibrium level of recruitment is

$$R_{eq} = \frac{\alpha\phi - \beta}{\phi}$$

The sustainable yield for a level of fishing mortality is $R_{eq} * YPR$, where YPR is the yield per recruit. MSY and F_{msy} are then obtained by finding the fishing mortality rate where yield is maximized, and this was accomplished by using the numerical Newton-Raphson technique to solve for the derivative of the yield curve.

Parameters Estimated Independently

The parameters estimated independently include the natural mortality (M) and survey catchability (q_{srv}). Most studies assume $M = 0.20$ for these species on the basis of their longevity. Fish from both sexes have frequently been aged as high as 25 years from samples collected during the annual trawl surveys. Zhang (1987) determined that the natural mortality rate for Alaska plaice is variable by sex and may range from 0.195 for males to 0.27 for females. Natural mortality was fixed at 0.25 for this assessment from the result of a previous assessment (Wilderbuer and Walters 1997, Table 8.1) where M was profiled over a range of values to explore the effect it has on the overall model fit and to the individual data components. The survey catchability was fixed at 1.0.

Parameters Estimated Conditionally

Parameter estimation is facilitated by comparing the model output to several observed quantities, such as the age compositions of the fishery and survey catches, the survey biomass, and the fishery catches. The general approach is to assume that deviations between model estimates and observed quantities are attributable to observation error and can be described with statistical distributions. Each data component provides a contribution to a total log-likelihood function, and parameter values that maximize the log-likelihood are selected.

The log-likelihoods of the age compositions were modeled with a multinomial distribution. The log of the multinomial function (excluding constant terms) is

$$n \sum_{t,a} p_{t,a} \ln(\hat{p}_{t,a})$$

where n_t is the number of fish aged, and p and \hat{p} are the observed and estimated age proportion at age.

The log-likelihood of the survey biomass was modeled with a lognormal distribution:

$$\lambda_2 \sum_t (\ln(obs_biom_t) - \ln(pred_biom_t))^2 / 2 * cv(t)^2$$

where obs_biom_t and $pred_biom_t$ are the observed and predicted survey biomass at time t , $cv(t)$ is the coefficient of variation of observed biomass in year t , and λ_2 is a weighting factor.

The predicted survey biomass for a given year is

$$q_{-srv} * \sum_a selsrv_a (\bar{N}_a * wt_a)$$

where $selsrv_a$ is the survey selectivity at age and wt_a is the population weight at age.

The log-likelihood of the catch biomass were modeled with a lognormal distribution:

$$\lambda_3 \sum_t (\ln(obs_cat_t) - \ln(pred_cat_t))^2$$

where obs_cat_t and $pred_cat_t$ are the observed and predicted catch. Because the catch biomass is generally thought to be observed with higher precision than other variables, λ_3 is given a very high value (hence low variance in the total catch estimate) so as to fit the catch biomass nearly exactly. This can be accomplished by varying the F levels, and the deviations in F are not included in the overall likelihood function. The overall likelihood function (excluding the catch component) is

$$\lambda_1 \left(\sum_t \varepsilon_t + \sum_a \gamma_a \right) + n \sum_{t,a} p_{t,a} \ln(\hat{p}_{t,a}) + \lambda_2 \sum_t (\ln(obs_biom_t) - \ln(pred_biom_t))^2 / 2 * cv(t)^2$$

For the model run in this analysis, λ_1 , λ_2 , and λ_3 were assigned weights of 1,1, and 500, respectively. The value for age composition sample size, n , was set to 200. The likelihood function was maximized by varying the following parameters:

Parameter type	Number
1) fishing mortality mean (μ)	1
2) fishing mortality deviations (ε_t)	33
3) recruitment mean (ν)	1
4) recruitment deviations (ν)	33
5) initial year mean (<i>meaninit</i>)	1
6) initial year deviations (γ)	22
7) fishery selectivity patterns	2
8) survey selectivity patterns	2
9) stock recruitment parameters	2
Total parameters	97

Finally, a Monte Carlo Markov Chain (MCMC) algorithm was used to obtain estimates of parameter uncertainty (Gelman et al. 1995). One million MCMC simulations were conducted, with every 1,000th sample saved for the sample from the posterior distribution. Ninety-five percent confidence intervals were produced as the values corresponding to the 5th and 95th percentiles of the MCMC evaluation. For this assessment, confidence intervals on total biomass and recruitment strength are presented.

Model Results

Substantial differences exist in the estimates of stock productivity and F_{msy} between model forms. When using the post-1977 year classes, the Ricker model estimates an F_{msy} of 0.20, which is substantially below the estimated $F_{40\%}$ of 0.59 (Table 9.9, Figure 9.6). When the

Beverton-Holt curve is used the stock-recruitment model is essentially a horizontal line through the data (Figure 9.7), as the steepness parameter is at its upper bound of 1.0. Both the Ricker and Beverton-Holt curves produce similar fits to the post-1989 year class data, but there is only a sparse amount of data in these later years to which a curve can be fit (Figures 9.8 and 9.9). Both curves estimate that productivity of Alaska plaice is so low that fishing at any level could not be sustained. Also note that the estimates of recruitment in the very last few years differ between the model fits. These recruitments represent cohorts that have yet to appear in any substantial numbers in the fishery and survey data, and thus have very little information to determine their magnitude. Given the uncertainties regarding which subset of years best characterize the current state of stock productivity, and the high degree to which the productivity estimates depend on this factor, it is not recommended that estimates of F_{msy} be used for management advice. The fitting of a stock-recruitment curve within the model remains a useful feature, and the following results are based upon the model that used a Ricker model fit to all available year classes.

The model results show that estimated total Alaska plaice biomass (ages 3+) increased from 1.1 million t in 1975 to a peak of 1.72 in 1982 (Figure 9.10, Table 9.10). Beginning in 1984, estimated total biomass declined to 1.06 million t in 2000 but has since increased to 1.74 million t in 2007 and is projected at 1.85 million t in 2008. The estimated survey biomass also shows a rapid increase to a peak biomass of 744,281 t in 1985, and a subsequent decline to a lower stable since then (Figure 9.11). The recent increase is the result of above average year classes spawned in 1998 and 2000 which are now at or nearing the age of maturity. The female spawning biomass trend is similar to the total biomass trend with a peak level estimated in 1985 and a slow decline thereafter until 2005 after which the spawning stock is estimated to be increasing.

Past assessments have estimated $F_{40\%}$ and $F_{35\%}$ at high levels for Alaska plaice (0.77 and 1.08, respectively). This is in part a result of the estimate of the fishery selectivity curve which indicated that Alaska plaice were 50% selected at an age of 10.9 years. However, these fishing mortality reference point estimates are quite high compared to other Bering Sea flatfish species and are computed from data collected in fisheries where Alaska plaice were not the fisheries target (85-87% of Alaska plaice are caught in the yellowfin sole fishery). For this assessment, fitting these fishery observations was de-emphasized by lowering the input sample sizes from 200 to 50. This had the effect of producing estimates of $F_{40\%}$ and $F_{35\%}$ at 0.59 and 0.81, respectively, and lowered the estimate of fishery selectivity to 10.4 years (Figure 9.2). The fits to the trawl survey age and length compositions are shown in Figures 9.13 and 9.14 and the fit to the fishery age and length compositions are shown in Figures 9.15 and 9.16.

The changes in stock biomass are primarily a function of recruitment variability, as fishing pressure has been relatively light. The fully selected fishing mortality estimates show a maximum value of 0.09 in 1988, and have averaged 0.03 from 1975-2007 (Figure 9.17). Estimated age-3 recruitment indicates high levels from the 1971-1980 year classes, averaging 1.9×10^9 (Figure 9.18, Table 9.10). From the 1981-1997 year classes, estimated recruitment has declined, averaging 1.1×10^9 . Recruitment is estimated to be improving since 1997 with above average strength recruitment from the 1998 and 2000 year classes. The 2002 year class may also be above average but have only been observed as three year olds in the 2005 and 2006 survey age samples. Although no age information is presently available from 2007, the survey length composition indicates the presence of small fish ranging from 10-20 cm which the model

estimates as a large incoming year class from 2003. This should be viewed with caution as it is a single observation of small fish which are only partially selected by the trawl (the model estimate has a large variance in Fig 9.18).

Projections and Harvest Alternatives

The reference fishing mortality rate for Alaska plaice is determined by the amount of reliable population information available (Amendment 56 of the Fishery Management Plan for the groundfish fishery of the Bering Sea/Aleutian Islands). Estimates of $B_{40\%}$, $F_{40\%}$, and $SPR_{40\%}$ were obtained from a spawner-per-recruit analysis. Assuming that the average recruitment from 1977-2003 year classes estimated in this assessment represents a reliable estimate of equilibrium recruitment, then an estimate of $B_{40\%}$ is calculated as the product of $SPR_{40\%}$ * equilibrium recruits, and this quantity is 144,500 t. The 2008 spawning biomass is estimated at 335,900 t. Since reliable estimates of 2008 spawning biomass (B), $B_{40\%}$, $F_{40\%}$, and $F_{35\%}$ exist and $B > B_{40\%}$ (335,900 t > 144,500 t), Alaska plaice reference fishing mortality is defined in tier 3a of Amendment 56. For this tier, F_{ABC} is constrained to be $\leq F_{40\%}$, and F_{OFL} is defined as $F_{35\%}$. The values of these quantities are:

2008 SSB estimate (B)	=	335,900 t
$B_{40\%}$	=	144,500 t
$F_{40\%}$	=	0.59
F_{ABC}	=	0.59
$F_{35\%}$	=	0.81
F_{OFL}	=	0.81

The estimated catch level for year 2008 associated with the overfishing level of $F = 0.81$ is 247,500 t. **The year 2008 recommended ABC associated with F_{ABC} of 0.59 is 194,100 t.** Projections of Alaska plaice female spawning biomass (described below) at a harvest rate equal to the average fishing mortality rate of the past five years indicate that the stock will increase over the next five years (Fig. 9.19).

A standard set of projections is required for each stock managed under Tiers 1, 2, or 3 of Amendment 56. This set of projections encompasses seven harvest scenarios designed to satisfy the requirements of Amendment 56, the National Environmental Policy Act, and the Magnuson-Stevens Fishery Conservation and Management Act (MSFCMA).

For each scenario, the projections begin with the vector of 2007 numbers at age estimated in the assessment. This vector is then projected forward to the beginning of 2008 using the schedules of natural mortality and selectivity described in the assessment and the best available estimate of total (year-end) catch for 2007. In each subsequent year, the fishing mortality rate is prescribed on the basis of the spawning biomass in that year and the respective harvest scenario. In each year, recruitment is drawn from an inverse Gaussian distribution whose parameters consist of maximum likelihood estimates determined from recruitments estimated in the assessment. Spawning biomass is computed in each year based on the time of peak spawning and the maturity and weight schedules described in the assessment. Total catch is assumed to equal the catch associated with the respective harvest scenario in all years. This projection scheme is run 1000 times to obtain distributions of possible future stock sizes, fishing mortality rates, and catches.

Five of the seven standard scenarios will be used in an Environmental Assessment prepared in conjunction with the final SAFE. These five scenarios, which are designed to provide a range of harvest alternatives that are likely to bracket the final TAC for 2008, are as follows (“ $max F_{ABC}$ ” refers to the maximum permissible value of F_{ABC} under Amendment 56):

Scenario 1: In all future years, F is set equal to $max F_{ABC}$. (Rationale: Historically, TAC has been constrained by ABC, so this scenario provides a likely upper limit on future TACs.)

Scenario 2: In all future years, F is set equal to a constant fraction of $max F_{ABC}$, where this fraction is equal to the ratio of the F_{ABC} value for 2008 recommended in the assessment to the $max F_{ABC}$ for 2008. (Rationale: When F_{ABC} is set at a value below $max F_{ABC}$, it is often set at the value recommended in the stock assessment.)

Scenario 3: In all future years, F is set equal to 50% of $max F_{ABC}$. (Rationale: This scenario provides a likely lower bound on F_{ABC} that still allows future harvest rates to be adjusted downward when stocks fall below reference levels.)

Scenario 4: In all future years, F is set equal to the 2003-2007 average F . (Rationale: For some stocks, TAC can be well below ABC, and recent average F may provide a better indicator of F_{TAC} than F_{ABC} .)

Scenario 5: In all future years, F is set equal to zero. (Rationale: In extreme cases, TAC may be set at a level close to zero.)

The recommended F_{ABC} and the maximum F_{ABC} are equivalent in this assessment, and five-year projections of the mean Alaska plaice harvest and spawning stock biomass for the remaining four scenarios are shown in Table 9.11.

Two other scenarios are needed to satisfy the MSFCMA’s requirement to determine whether the Alaska plaice stock is currently in an overfished condition or is approaching an overfished condition. These two scenarios are as follows (for Tier 3 stocks, the MSY level is defined as $B_{35\%}$):

Scenario 6: In all future years, F is set equal to F_{OFL} . (Rationale: This scenario determines whether a stock is overfished. If the stock is expected to be above its MSY level in 2008 under this scenario, then the stock is not overfished.)

Scenario 7: In 2008 and 2009, F is set equal to $max F_{ABC}$, and in all subsequent years, F is set equal to F_{OFL} . (Rationale: This scenario determines whether a stock is approaching an overfished condition. If the stock is expected to be above its MSY level in 2010 under this scenario, then the stock is not approaching an overfished condition.)

The results of these two scenarios indicate that Alaska plaice are neither overfished nor approaching an overfished condition. With regard to assessing the current stock level, the

expected stock size in the year 2008 of scenario 6 is more than twice its $B_{35\%}$ value of 126,400 t. With regard to whether the stock is likely to be in an overfished condition in the near future, the expected stock size in the year 2010 of scenario 7 is also greater than its $B_{35\%}$ value.

Scenario Projections and Two-Year Ahead Overfishing Level

In addition to the seven standard harvest scenarios, Amendments 48/48 to the BSAI and GOA Groundfish Fishery Management Plans require projections of the likely OFL two years into the future. While Scenario 6 gives the best estimate of OFL for 2008, it does not provide the best estimate of OFL for 2009, because the mean 2009 catch under Scenario 6 is predicated on the 2008 catch being equal to the 2008 OFL, whereas the actual 2008 catch will likely be less than the 2008 ABC. Therefore, the projection model was re-run with the 2008 catch fixed equal to the 2007 catch and the 2009 fishing mortality rate fixed at F_{ABC} .

Year	Catch	ABC	OFL
2008	19,176	194,100	247,500
2009	19,176	216,600	276,700

Ecosystem considerations

Ecosystem Effects on the stock

1) Prey availability/abundance trends

The feeding habits of juvenile Alaska plaice are relatively unknown, although the larvae are relatively large at hatching (5.85 mm) with more advanced development than other flatfish (Pertseva-Ostroumova 1961).

For adult fish, Zhang (1987) found that the diet consisted primarily of polychaetes and amphipods regardless of size. For fish under 30 cm, polychaetes contributed 63% of the total diet with sipunculids (marine worms) and amphipods contributing 21.7% and 11.6%, respectively. For fish over 30 cm, polychaetes contributed 75.2% of the total diet with amphipods and echiurans (marine worms) contributing 6.7% and 5.7%, respectively. Similar results were in stomach sampling from 1993-1996, with polychaetes and marine worms composing the majority of the Alaska plaice diet (Lang et al. 2003). McConnaughey and Smith (2000) contrasted the food habits of several flatfish between areas of high and low CPUE, using aggregated data from 1982 to 1994. For Alaska plaice, the diets were nearly identical with 76.5% of the diet composed of polychaetes and unsegmented coelomate worms in the high CPUE areas as compared to 83.1% in the low CPUE areas.

2) Predator population trends

Alaska plaice contribute a relatively small portion of the diets of Pacific cod, Pacific halibut, and yellowfin sole as compared with other flatfish. Total consumption estimates of Alaska plaice from 1993 to 1996 ranged from 0 t in 1996 to 574 t in 1994 (Lang et al. 2003). Consumption by

yellowfin sole is upon fish < 2 cm whereas consumption by Pacific halibut is upon fish > 19 cm (Lang et al. 2003).

3) Changes in habitat quality

The habitats occupied by Alaska plaice are influenced by temperature, which has shown considerable variation in the eastern Bering Sea in recent years. For example, the timing of spawning and advection to nursery areas are expected to be affected by environmental variation. Musienko (1970) reported that spawning occurs immediately after the ice melt, with peak spawning occurring at water temperatures from -1.53 to 4.11. In 1999, one of the coldest years in the eastern Bering Sea, the distribution was shifted further to the southeast than it was during 1998-2002. However, in 2003, one of the warmest years in the EBS, the distribution was shifted further to the southeast than observed in 1999.

Fishery effects on the ecosystem

Alaska plaice are not a targeted species and are harvested in a variety of fisheries in the BSAI area. Since 2002, when single-species management for Alaska plaice was initiated, harvest estimates by fishery are available. Most Alaska plaice are harvested within the yellowfin sole fishery, accounting for 81% - 87% of the catch in 2002-2006. Flathead sole, rock sole, and Pacific cod fisheries make up the remainder of the catch. The ecosystem effects of the yellowfin sole fishery can be found with the yellowfin sole assessment in this SAFE document.

Due to the minimal consumption estimates of Alaska plaice (Lang et al. 2003) by other groundfish predators, the yellowfin sole fishery does not have a significant impact upon those species preying upon Alaska plaice. Additionally, the relatively light fishing mortality rates experienced by Alaska plaice are not expected to have significant impacts on the size structure of the population or the maturity and fecundity at age. It is not known what effects the fishery may have on the maturity-at-age of Alaska plaice. The yellowfin sole fishery, however, does contribute substantially to the total discards in the EBS, as indicated by the discarding of Alaska plaice discussed in this assessment, and general discards within this fishery discussed in the yellowfin sole assessment.

Summary

In summary, several quantities pertinent to the management of the Alaska plaice are listed below.

<u>Quantity</u>	<u>Value</u>
M	0.25
Tier	3a
Year 2008 Total Biomass	1,854,000 t
Year 2008 Spawning stock biomass	335,900 t
$B_{100\%}$	361,100 t
$B_{40\%}$	144,500 t
$B_{35\%}$	126,400 t
F_{OFL}	0.81
Maximum F_{ABC}	0.59
Recommended F_{ABC}	0.59
OFL	247,500 t
Maximum allowable ABC	194,100 t
<u>Recommended ABC</u>	<u>194,100 t</u>

References

- Gelman, A., J.B. Carlin, H.S. Stern, and D.A. Rubin. 1995. Bayesian data analysis. Chapman and Hall, New York. 552 pp.
- Hilborn, R. and C.J. Walters. 1992. Quantitative fisheries stock assessment: choices, dynamics, and uncertainty. Chapman and Hall, New York. 570 pp.
- Haflinger, K. 1981. A survey of benthic infaunal communities of the southeastern Bering Sea shelf. In D.W Hood and J.A. Calder (eds), The eastern Bering Sea shelf: oceanography and resources. Univ. of Wash. Press, Seattle, pp 1091-1104.
- Kappenman, R. F. 1992. Estimation of the fishing power correction factor. Processed Report 92-01, 10 p. Alaska Fish. Sci. Center, Natl. Mar. Fish. Serv., NOAA, 7600 Sand Point Way NE, Seattle, WA 98115.
- Kimura, D.K. and S. Chikuni. 1987. Mixtures of empirical distributions: An iterative application of the age-length key. Biometrics 43:23-34.
- Lang, G.M., C.W. Derah, and P.A. Livingston. 2003. Groundfish food habits and predation on commercially important prey species in the eastern Bering Sea from 1993 to 1996. U.S. Dep. Commer., AFSC Proc. Rep. 2003-04. 351 pp.

- McConnaughey, R.A. and K.R. Smith. 2000. Associations between flatfish abundance and surficial sediments in the eastern Bering Sea. *Can J. Fish. Aquat. Sci.* 2410-2419.
- Musienko, L.N. 1970. Reproduction and development of Bering Sea fishes. Tr. Vses. Nauchno-issled. Inst. Morsk. Rybn. Khoz. Okeanogr. 70 (Izv. Tikhookean. Nauchno-issled. Inst. Morsk. Rybn. Khoz. Okeanogr. 72:166-224) Transl. *In Sov. Fish. Invest. Northeast Pac.*, pt. V:161-224. *Isr. Program Sci. Transl.*, 1972. Avail. From
- Pertseva-Ostroumova. 1961. The reproduction and development of far eastern flounders. *Akad. Nauk SSSR Inst. Okeanologii*, 484 p. (Transl. by Fish. Res. Bd. Can., 1967, Transl. Ser. 856, 1003 p.)
- Walters, G. E., and T. K. Wilderbuer. 1990. Other flatfish. In Stock Assessment and Fishery Evaluation Document for Groundfish Resources in the Bering Sea/Aleutian Islands Region as Projected for 1991, p.129-141. North Pacific Fishery Management Council, P.O. Box 103136, Anchorage Alaska 99510.
- Wilderbuer, T. K., and G. E. Walters. 1997. Other flatfish. In Stock Assessment and Fishery Evaluation Document for Groundfish Resources in the Bering Sea/Aleutian Islands Region as Projected for 1998, p.271-296. North Pacific Fishery Management Council, P.O. Box 103136, Anchorage Alaska 99510.
- Zhang, C. I. 1987. Biology and population dynamics of Alaska plaice, *Pleuronectes quadrituberculatus*, in the eastern Bering Sea. Ph. D. dissertation, University of Washington:1-225.
- Zhang, C. I., T.K. Wilderbuer, and G.E. Walters. 1998. Biological characteristics and fishery assessment of Alaska plaice, *Pleuronectes quadrituberculatus*, in the Eastern Bering Sea. *Marine Fisheries Review* 60(4), 16-27.

Table 9.1. Harvest (t) of Alaska plaice from 1977-2007

<u>Year</u>	<u>Harvest</u>
1977	2589
1978	10420
1979	13672
1980	6902
1981	8653
1982	6811
1983	10766
1984	18982
1985	24888
1986	46519
1987	18567
1988	61638
1989	14134
1990	10926
1991	15003
1992	18074
1993	13846
1994	10882
1995	19172
1996	16096
1997	21236
1998	14296
1999	13997
2000	14487
2001	8685
2002	12176
2003	9978
2004	7572
2005	11079
2006*	17202
2007*	19176

*NMFS Regional Office Report through Sept 8, 2007

Table 9.2. Research catches (t) of Alaska plaice in the BSAI area from 1977 to 2005.

Year	Research Catch (t)
1977	4.28
1978	4.94
1979	17.15
1980	12.02
1981	14.31
1982	26.77
1983	43.27
1984	32.42
1985	23.24
1986	19.66
1987	19.74
1988	39.42
1989	31.10
1990	32.29
1991	29.79
1992	15.14
1993	19.71
1994	22.48
1995	28.47
1996	18.26
1997	22.59
1998	17.17
1999	18.95
2000	15.98
2001	20.45
2002	15.07
2003	15.39
2004	18.03
2005	22.52
2006	28.50
2007	18.80

Table 9.3. Restrictions on the “other flatfish” fishery from 1995 to 2007 in the Bering Sea – Aleutian Islands management area. Unless otherwise indicated, the closures were applied to the entire BSAI management area. Zone 1 consists of areas 508, 509, 512, and 516, whereas zone 2 consists of areas 513, 517, and 521.

Year	Dates	Bycatch Closure
1995	2/21 – 3/30	First Seasonal halibut cap
	4/17 – 7/1	Second seasonal halibut cap
	8/1 – 12/31	Annual halibut allowance
1996	2/26 – 4/1	First Seasonal halibut cap
	4/13 – 7/1	Second seasonal halibut cap
	7/31 – 12/31	Annual halibut allowance
1997	2/20 – 4/1	First Seasonal halibut cap
	4/12 – 7/1	Second seasonal halibut cap
	7/25 – 12/31	Annual halibut allowance
1998	3/5 – 3/30	First Seasonal halibut cap
	4/21 – 7/1	Second seasonal halibut cap
	8/16 – 12/31	Annual halibut allowance
1999	2/26 – 3/30	First Seasonal halibut cap
	4/27 – 7/04	Second seasonal halibut cap
	8/31 – 12/31	Annual halibut allowance
2000	¾ – 3/31	First Seasonal halibut cap
	4/30 – 7/03	Second seasonal halibut cap
	8/25 – 12/31	Annual halibut allowance
2001	3/20 – 3/31	First Seasonal halibut cap
	4/27 – 7/01	Second seasonal halibut cap
	8/24 – 12/31	Annual halibut allowance
2002	2/22 – 12/31	Red King crab cap (Zone 1 closed)
	3/1 – 3/31	First Seasonal halibut cap
	4/20 – 6/29	Second seasonal halibut cap
	7/29 – 12/31	Annual halibut allowance
2003	2/18 – 3/31	First Seasonal halibut cap
	4/1 – 6/21	Second seasonal halibut cap
	7/31 – 12/31	Annual halibut allowance
2004	2/24 – 3/31	First Seasonal halibut cap
	4/10 – 12/31	Bycatch status
2005	3/1 - 3/31	First Seasonal halibut cap
	4/22-6/30	Second Seasonal halibut cap
	5/9-12/31	Bycatch status, TAC attained
2006	2/21 - 3/31	First Seasonal halibut cap
	4/5 – 12/31	Red King crab cap (Zone 1 closed)
	4/12 – 5/31	Second seasonal halibut cap
	5/26	TAC attained, 7,000 t reserve released
	8/7 – 12/31	Annual halibut allowance
2007	2/17-3/31	First seasonal halibut cap
	4/1-6/21	Second seasonal halibut cap
	7/31-12/31	Annual halibut allowance

Table 9.4 Discarded and retained BSAI Alaska plaice catch (t) for 2002-2004, from NMFS Alaska regional office 'blend' (2002) and catch accounting system (2003 - 2005) data.

year	Discard	Retained	Total	Percent discarded
2003	11806	370	12176	0.97
2003	9428	350	9778	0.96
2004	7193	379	7572	0.95
2005	10293	786	11079	0.93
2006	14746	2564	17310	0.85

Table 9.5. Alaska plaice sample sizes from the BSAI fishery. The hauls columns refer to the number of hauls where from which either lengths or read otoliths were obtained.

Year	Total hauls	Hauls with lengths	# lengths	hauls w/lengths	hauls w/otoliths	# otoliths collected	# aged
1982	334	152	14274	27	27	298	298
1983	353	118	11624				
1984	355	151	14026	32	457		
1985	358	168	10914	24	430		
1986	354	236	12349				
1987	360	174	8535				
1988	373	170	7079	10	10	284	284
1989	373	206	7717				
1990	371	215	7739	10	228		
1991	372	235	8163				
1992	356	219	7584	10	10	311	311
1993	375	241	8365	4	4	183	183
1994	376	249	9300	6	6	228	228
1995	376	252	9919	11	11	287	285
1996	375	254	10186	5	250		
1997	376	248	10143	3	82		
1998	375	281	10101	14	14	420	416
1999	373	268	13024	13	297		
2000	372	250	9803	16	16	368	359
2001	375	261	10990	16	16	339	335
2002	375	251	8409	24	24	359	355
2003	376	252	8343	15	320		
2004	375	262	8578	17	325		
2005	373	262	9284	20	20	341	337
2006	376	255	12097	18	18	368	362
2007	376	261	11729	43	343		

Table 9.6. Estimated biomass and standard deviations (t) of Alaska plaice from the eastern Bering Sea trawl survey.

<u>Year</u>	<u>Biomass estimate</u>	<u>Standard Deviation</u>
1975	103,500	11,600
1979	277,200	31,100
1980	354,000	39,800
1981	535,800	60,200
1982	715,400	64,800
1983	743,000	65,100
1984	789,200	35,800
1985	580,000	61,000
1986	553,900	63,000
1987	564,400	57,500
1988	699,400	140,000
1989	534,000	58,800
1990	522,800	50,000
1991	529,000	50,100
1992	530,400	56,400
1993	515,200	50,500
1994	623,100	53,300
1995	552,292	62,600
1996	529,300	67,500
1997	643,400	73,200
1998	452,600	58,700
1999	546,522	47,000
2000	443,620	67,600
2001	540,458	68,600
2002	428,519	53,800
2003	467,326	97,400
2004	488,217	63,800
2005	503,861	55,698
2006	636,971	81,547
2007	421,765	37,831

Table 9.7. Alaska plaice population numbers at age estimated from the NMFS eastern Bering Sea groundfish surveys and age readings of sampled fish.

Year	Number at age (millions)															
	3	4	5	6	7	8	9	10	11	12	13	14	15	16+		
1982	0.49	0.20	23.58	74.93	134.98	161.68	146.68	128.17	152.75	148.45	188.07	163.15	98.46	52.17		
1988	0.00	0.07	8.47	18.07	97.84	74.61	138.52	67.08	158.83	74.74	33.19	98.87	11.57	216.90		
1992	0.00	8.86	30.17	6.78	37.24	68.52	51.45	51.50	78.08	46.41	36.36	44.36	33.74	231.10		
1993	0.00	0.00	10.19	51.37	45.07	65.83	99.24	24.56	20.83	54.33	88.53	36.94	56.61	209.74		
1994	0.00	0.00	24.02	36.20	123.52	107.60	45.82	91.80	38.82	25.88	113.13	51.75	76.37	232.74		
1995	0.00	0.00	6.19	69.33	60.37	133.83	60.79	36.73	61.29	31.22	28.09	41.37	54.22	268.52		
1998	0.00	1.10	8.77	31.04	77.79	75.16	105.41	53.12	60.67	64.33	29.41	42.91	32.07	150.46		
2000	0.00	0.13	10.67	5.68	44.75	53.88	135.66	75.86	67.11	44.94	40.88	32.04	17.02	258.41		
2001	0.00	0.00	6.35	27.96	24.46	124.33	68.93	174.05	57.51	93.86	34.35	67.23	14.35	252.29		
2002	0.00	0.94	3.72	30.78	42.49	36.86	74.11	58.01	79.95	35.25	56.29	23.99	48.05	178.14		
2005	1.50	6.26	23.51	55.63	57.86	82.33	45.88	71.56	42.7	61.25	74.12	41.24	32.7	152.17		
2006	0.35	14.33	100.69	52.31	122.05	128.35	93.90	56.38	43.66	45.57	50.4	78.92	60.76	152.9		

Table 9.8. Alaska plaice sample sizes from the BSAI trawl survey. The hauls columns refer to the number of hauls from which either lengths or aged otoliths were obtained.

Year	Total hauls	Hauls with lengths	# lengths	hauls w/lengths	hauls w/otoliths	# otoliths collected	# aged
1982	334	152	14274	27	27	298	298
1983	353	118	11624				
1984	355	151	14026	32	457		
1985	358	168	10914	24	430		
1986	354	236	12349				
1987	360	174	8535				
1988	373	170	7079	10	10	284	284
1989	373	206	7717				
1990	371	215	7739	10	228		
1991	372	235	8163				
1992	356	219	7584	10	10	311	311
1993	375	241	8365	4	4	183	183
1994	376	249	9300	6	6	228	228
1995	376	252	9919	11	11	287	285
1996	375	254	10186	5	250		
1997	376	248	10143	3	82		
1998	375	281	10101	14	14	420	416
1999	373	268	13024	13	297		
2000	372	250	9803	16	16	368	359
2001	375	261	10990	16	16	339	335
2002	375	251	8409	24	24	359	355
2003	376	252	8343	15	320		
2004	375	262	8578	17	325		
2005	373	262	9284	20	20	341	337
2006	376	255	12097	18	18	368	362
2007	376	261	11729	43	343		

Table 9.9. Estimates of management parameters associated with fitting the Ricker and Beverton-Holt stock recruitment relationships to two different time spans of data, with standard deviations in parentheses. Standard deviations were not obtained for the case of fitting the Beverton-Holt model to year classes 1989-2001 because the Hessian was not positive definite.

SR model	year classes	F_{40}	F_{msv}	$B_{msv}(t)$	MSY (t)	Notes
Ricker	77-01	0.76 (0.05)	0.20 (0.18)	135460 (14249)	29174 (22198)	
Ricker	89-01	0.75 (0.05)	0.0003 (0.008)	1271.7 (29070)	1.0 (27.62)	
Beverton-Holt	77-01	0.76 (0.05)	21.9 (53.92)	21025 (34821)	84320 (13632)	Steepness at upper bound of 1.0
Beverton-Holt	89-01	0.75 (0.05)	3.83×10^{-7}	1.0	6.19×10^{-7}	Hessian not positive definite, steepness at lower bound of 0.2

Table 9.10. Estimated total biomass (ages 3+), female spawner biomass, and recruitment (age 3), with comparison to the 2005 SAFE estimates.

	Female spawning biomass		Total biomass (t)		Age 3 recruitment (millions)	
	2007	2006	2007	2006	2007	2006
1975	166,753	167,475	1,124,670	1,128,280	2,075	2,088
1976	202,247	203,153	1,279,740	1,283,880	3,603	3,618
1977	240,262	241,290	1,412,120	1,416,460	1,911	1,909
1978	272,972	274,006	1,523,380	1,527,820	1,882	1,882
1979	298,103	299,115	1,611,530	1,615,620	1,811	1,803
1980	326,633	327,668	1,669,960	1,673,500	1,311	1,306
1981	358,743	359,869	1,701,840	1,704,800	1,476	1,475
1982	398,742	399,965	1,712,070	1,714,480	1,425	1,425
1983	429,816	431,048	1,703,810	1,705,740	1,507	1,508
1984	454,864	456,023	1,651,470	1,652,860	678	674
1985	460,944	461,885	1,574,490	1,575,640	744	749
1986	452,213	452,926	1,482,390	1,483,630	1,318	1,327
1987	436,701	437,197	1,410,710	1,412,540	849	866
1988	416,588	416,912	1,304,420	1,307,230	1,090	1,108
1989	390,171	390,395	1,274,980	1,279,340	1,686	1,712
1990	373,705	373,909	1,240,970	1,247,240	870	896
1991	356,317	356,619	1,216,040	1,224,390	1,285	1,310
1992	337,118	337,628	1,182,840	1,194,200	874	926
1993	320,470	321,366	1,170,990	1,186,240	1,453	1,517
1994	311,170	312,579	1,157,730	1,175,330	1,009	1,007
1995	305,476	307,523	1,139,370	1,162,370	1,114	1,239
1996	297,628	300,420	1,110,660	1,138,440	634	690
1997	294,406	298,113	1,074,860	1,111,700	832	1,025
1998	287,679	292,356	1,041,050	1,088,170	760	912
1999	287,149	293,020	1,014,590	1,071,720	1,068	1,180
2000	282,069	289,084	1,005,860	1,064,520	1,457	1,321
2001	278,535	287,616	1,035,200	1,085,510	2,012	1,760
2002	270,704	281,939	1,071,110	1,112,580	1,586	1,566
2003	262,809	277,394	1,163,310	1,146,440	2,942	1,431
2004	255,944	272,757	1,261,550	1,165,620	2,045	857
2005	256,635	274,126	1,369,590	1,253,650	2,295	3,513
2006	262,449	277,725	1,589,960	1,297,570	5,860	996
2007	278,876		1,741,800		1,524	

Table 9.11. Projections of spawning biomass (t), catch, fishing mortality rate, and catch (t) for each of the several scenarios. The values of $B_{40\%}$ and $B_{35\%}$ are 144,500 t and 126,400 t, respectively. ABC is highlighted.

Scenarios 1 and 2

Maximum ABC harvest permissible

Female

Year	spwn bio	catch	F
2007	301.762	19.176	0.053
2008	304.658	194.100	0.591
2009	282.647	158.156	0.591
2010	288.46	147.063	0.591
2011	307.783	148.360	0.591
2012	304.114	158.608	0.591
2013	289.939	174.665	0.591
2014	246.559	168.503	0.591
2015	204.898	137.876	0.591
2016	176.198	110.091	0.590
2017	160.78	92.957	0.580
2018	153.473	84.348	0.571
2019	150.185	80.335	0.567
2020	148.887	78.469	0.565

Scenario 3

1/2 Maximum ABC harvest permissible

Female

Year	spwn bio	catch	F
2007	301.762	19.176	0.053
2008	322.904	97.050	0.264
2009	316.01	190.245	0.591
2010	306.292	164.741	0.591
2011	316.909	157.634	0.591
2012	308.63	163.215	0.591
2013	292.117	176.733	0.591
2014	247.587	169.058	0.591
2015	205.374	137.282	0.591
2016	176.414	108.849	0.590
2017	160.874	92.568	0.580
2018	153.511	84.140	0.572
2019	150.2	80.035	0.567
2020	148.893	77.994	0.565

Scenario 4

Harvest at average F over the past 5 years

Female

Year	spwn bio	catch	F
2007	301.762	19.176	0.053
2008	336.478	15.465	0.039
2009	380.102	15.643	0.035
2010	431.947	17.745	0.035
2011	487.344	20.105	0.035
2012	515.751	22.697	0.035
2013	535.44	25.707	0.035
2014	524.465	27.838	0.035
2015	503.562	27.769	0.035
2016	475.779	26.353	0.035
2017	448.947	24.643	0.035
2018	425.467	23.052	0.035
2019	405.409	21.676	0.035
2020	388.352	20.524	0.035

Scenario 5

No fishing

Female

Year	spwn bio	catch	F
2007	301.762	19.176	0
2008	338.901	0	0
2009	388.975	0	0
2010	446.855	0	0
2011	508.363	0	0
2012	543.05	0	0
2013	569.354	0	0
2014	565.196	0	0
2015	550.745	0	0
2016	527.839	0	0
2017	504.101	0	0
2018	482.218	0	0
2019	462.591	0	0
2020	445.03	0	0

Table 9.11- continued

Scenario 6
Determination of
overfishing

B35=126.4

Year	Female		
	spwn bio	catch	F
2007	301.762	19.176	0.053
2008	293.383	247.508	0.809
2009	256.782	181.236	0.809
2010	258.284	161.869	0.809
2011	275.815	161.839	0.809
2012	270.362	174.505	0.809
2013	252.495	192.371	0.809
2014	206.537	177.448	0.809
2015	167.304	136.474	0.806
2016	145.459	103.368	0.765
2017	136.837	88.837	0.737
2018	133.973	83.927	0.727
2019	133.114	82.236	0.723
2020	133.042	81.694	0.722

Scenario 7
Determination of whether Alaska plaice are
approaching
an overfished condition

Year	Female		
	spwn bio	catch	F
2007	301.762	19.176	0.053
2008	304.658	194.097	0.591
2009	282.648	158.157	0.591
2010	280.494	189.566	0.809
2011	286.238	175.338	0.809
2012	275.017	180.687	0.809
2013	254.495	195.068	0.809
2014	207.364	178.589	0.809
2015	167.627	136.955	0.806
2016	145.573	103.563	0.765
2017	136.871	88.897	0.737
2018	133.981	83.943	0.727
2019	133.115	82.239	0.723
2020	133.042	81.694	0.722







Figure 9.1 Locations of Alaska plaice catch in 2006, by month. The harvest primarily occurred in the yellowfin sole fishery and rock sole fisheries.









Figure 9.2 Estimated survey biomass and 95% confidence intervals from NMFS eastern Bering Sea bottom trawl surveys.




Figure 9.3. Normalized residuals of Alaska plaice survey biomass (from lowess fit; solid line) and average temperature (dashed line)




Figure 9.4. Mean length of Alaska plaice for ages 5-12, by year, from survey sampling




Figure 9.5. Estimated weight-at-age relationship used in the 2006 assessment.




Figure 9.6. Estimated Ricker stock recruitment relationship for Alaska plaice using the year classes 1977–2001, with the replacement lines for $F_{40\%}$ (dashed line) and no fishing (dotted line).




Figure 9.7. Estimated Beverton-Holt stock recruitment relationship using for Alaska plaice using the year classes 1975 –2001, with the replacement lines for $F_{40\%}$ (dashed line) and no fishing (dotted line).




Figure 9.8. Estimated Ricker stock recruitment relationship using for Alaska plaice using the year classes 1989 –2001, with the replacement lines for F40% (dashed line) and no fishing (dotted line).




Figure 9.9. Estimated Beverton-Holt stock recruitment relationship using for Alaska plaice using the year classes 1989 –2001, with the replacement lines for $F_{40\%}$ (dashed line) and no fishing (dotted line).




Figure 9.10 Estimated beginning year total biomass of Alaska plaice from the assessment model, with 95% confidence intervals from MCMC integration.




Figure 9.11--Observed (data points) and predicted (solid line) survey biomass of Alaska plaice.




Figure 9.12 Model estimates of survey and fishery selectivity.




Figure 9.13 Survey age composition (solid line = observed, dotted line = predicted).




Figure 9.14 Survey length composition by year (solid line = observed, dotted line = predicted)

fishery ages




Figure 9.15 Fishery age composition by year (solid line = observed, dotted line = predicted)




Figure 9.16 Fishery length composition by year (solid line = observed, dotted line = predicted)




Figure 9.17 Estimated fully selected fishing mortality.




Figure 9.18--Estimated recruitment (age 3) of Alaska plaice with 95% confidence intervals obtained from MCMC integration. (The error bar for the 2003 observation is off the chart).




Figure 9.19 Projection of Alaska plaice at the harvest rate of the average of the past five years.